

# SPRAVATO<sup>®</sup> Treatment Experience



Learn what to expect  
with **SPRAVATO<sup>®</sup>**  
treatment experience

## What is **SPRAVATO<sup>®</sup>** (esketamine) **III** nasal spray?

**SPRAVATO<sup>®</sup>** is a prescription medicine used:

- with or without an antidepressant taken by mouth, to treat adults with treatment-resistant depression (TRD)
- with an antidepressant taken by mouth, to treat depressive symptoms in adults with major depressive disorder (MDD) with suicidal thoughts or actions

**SPRAVATO<sup>®</sup>** is not for use as a medicine to prevent or relieve pain (anesthetic). It is not known if **SPRAVATO<sup>®</sup>** is safe or effective as an anesthetic medicine.

It is not known if **SPRAVATO<sup>®</sup>** is safe and effective for use in preventing suicide or in reducing suicidal thoughts or actions. **SPRAVATO<sup>®</sup>** is not for use in place of hospitalization if your healthcare provider determines that hospitalization is needed, even if improvement is experienced after the first dose of **SPRAVATO<sup>®</sup>**.

It is not known if **SPRAVATO<sup>®</sup>** is safe and effective in children.

## IMPORTANT SAFETY INFORMATION

### What is the most important information I should know about **SPRAVATO<sup>®</sup>**?

#### **SPRAVATO<sup>®</sup>** can cause serious side effects, including:

- **Sedation, dissociation, and respiratory depression.** **SPRAVATO<sup>®</sup>** may cause sleepiness (sedation), fainting, dizziness, spinning sensation, anxiety, or feeling disconnected from yourself, your thoughts, feelings, space and time (dissociation), breathing problems (respiratory depression and respiratory arrest)
  - Tell your healthcare provider right away if you feel like you cannot stay awake or if you feel like you are going to pass out.
  - Your healthcare provider must monitor you for serious side effects for at least 2 hours after taking **SPRAVATO<sup>®</sup>**. Your healthcare provider will decide when you are ready to leave the healthcare setting.

*(continued)*

Please see Important Safety Information throughout, full [Prescribing Information](#), including Boxed WARNINGS, and [Medication Guide](#) for **SPRAVATO<sup>®</sup>** and discuss any questions you may have with your healthcare provider.


# SPRAVATO<sup>®</sup> Treatment Experience

SPRAVATO<sup>®</sup> is not a pill you have to take every day. It is an FDA-approved nasal spray, taken in healthcare offices that are certified to administer SPRAVATO<sup>®</sup>. It can be taken with or without an antidepressant.

Here are the steps to get started on SPRAVATO<sup>®</sup> and what to expect:

## Consultation

Your first appointment will be a consultation. An HCP will review your medical history and treatment goals to determine if SPRAVATO<sup>®</sup> is right for you. Your HCP may offer SPRAVATO<sup>®</sup>, but if not, you and your provider can use [Find a Center](#) to find a provider who does.




## Insurance & Approval

Your provider's office will work with you to get insurance approval before starting treatment. After you are approved, SPRAVATO<sup>®</sup> will be delivered to the office. 9 in 10 commercially insured patients are covered.\* SPRAVATO<sup>®</sup> is also covered under Medicare and Medicaid.<sup>††</sup>



## First Month

You will take SPRAVATO<sup>®</sup> twice a week for four weeks. After that, you and your provider will determine if treatment should be continued and will have the flexibility to decide your treatment frequency.



## Monitoring

After each treatment, you will stay in the healthcare office for at least 2 hours, where a healthcare provider will monitor you for any serious side effects. Because of possible side effects affecting mental alertness and motor coordination, you won't be able to drive, operate machinery, or do anything where you need to be completely alert until the next day, following a restful sleep.



\*TRD data only. Managed Markets Insights and Technology, LLC<sup>™</sup>, a trademark of MMIT, as of September 2024. Collected as of September 2024 and may change. This percentage may not represent 100% of formulary lives due to data limitations.

<sup>††</sup>Coverage relevant to your patients may vary by payer and may change over time. Contact the plan for the most current policies.

## IMPORTANT SAFETY INFORMATION (continued)

- **Abuse and misuse.** There is a risk for abuse and misuse with SPRAVATO<sup>®</sup>, which may lead to physical and psychological dependence. Your healthcare provider should check you for signs of abuse, misuse, and dependence before and during treatment.
  - Tell your healthcare provider if you have ever abused or been dependent on alcohol, prescription medicines, or street drugs.
  - Your healthcare provider can tell you more about the differences between physical and psychological dependence in drug addiction.
- **SPRAVATO<sup>®</sup> Risk Evaluation and Mitigation Strategy (REMS).** Because of the risks for sedation, dissociation, respiratory depression, and abuse and misuse, SPRAVATO<sup>®</sup>

is only available through a restricted program called the SPRAVATO<sup>®</sup> Risk Evaluation and Mitigation Strategy (REMS) Program. SPRAVATO<sup>®</sup> can only be administered at healthcare settings certified in the SPRAVATO<sup>®</sup> REMS Program. Patients treated in outpatient healthcare settings (such as medical offices and clinics) must be enrolled in the program.

- **Increased risk of suicidal thoughts and actions.** Antidepressant medicines may increase suicidal thoughts and actions in some people 24 years of age and younger, **especially within the first few months of treatment or when the dose is changed. SPRAVATO<sup>®</sup> is not for use in children.**


(continued)

Please see Important Safety Information throughout, full [Prescribing Information](#), including **Boxed WARNINGS**, and [Medication Guide](#) for SPRAVATO<sup>®</sup> and discuss any questions you may have with your healthcare provider.

**Here are the steps to get started on SPRAVATO<sup>®</sup> and what to expect (continued):**

**Second Month & Beyond**

You will take SPRAVATO<sup>®</sup> once a week during the second month. After that, some people continue treatment once a week, while others might switch to every other week. Throughout your treatment, you and your healthcare provider will work together to decide if you should continue with SPRAVATO<sup>®</sup> and how often you should take it.



**Visit SPRAVATO<sup>®</sup> YouTube Page**

Visit SPRAVATO<sup>®</sup> YouTube to learn more about real patients experience with SPRAVATO<sup>®</sup>.

**Learn More**

**Begin To Feel Like Yourself Again**

SPRAVATO<sup>®</sup> works to rapidly reduce depressive symptoms. In short term clinical trials, SPRAVATO<sup>®</sup> showed reduction of depression symptoms at 4 weeks compared to placebo, with most of the improvement seen as soon as 24 hours after treatment.

The most common side effects are: feeling disconnected from yourself, your thoughts, feelings and things around you; dizziness; nausea; feeling sleepy; spinning sensation; decreased feeling of sensitivity (numbness); feeling anxious; lack of energy; increased blood pressure; vomiting; feeling drunk; headache; and feeling very happy or excited.

If these common side effects occur, they usually happen right after taking SPRAVATO<sup>®</sup> and go away the same day.



**IMPORTANT SAFETY INFORMATION (continued)**

◦ Depression and other serious mental illnesses are the most important causes of suicidal thoughts and actions. Some people may have a higher risk of having suicidal thoughts or actions. These include people who have (or have a family history of) depression or a history of suicidal thoughts or actions.

**• How can I watch for and try to prevent suicidal thoughts and actions in myself or a family member?**

- Pay close attention to any changes, especially sudden changes, in mood, behavior, thoughts, or feelings, or if you develop suicidal thoughts or actions.
- Tell your healthcare provider right away if you have any new or sudden changes in mood, behavior, thoughts, or feelings, or if you develop suicidal thoughts or actions.
- Keep all follow-up visits with your healthcare provider as scheduled. Call your healthcare provider between visits as needed, especially if you have concerns about symptoms.

**• Tell your healthcare provider or get emergency help right away if you or your family member have any of the following symptoms, especially if they are new, worse, or worry you:**

- |   |   |
|---|---|
| ◦ thoughts about suicide or dying                     | ◦ suicide attempts                          |
| ◦ new or worse depression                             | ◦ new or worse anxiety                      |
| ◦ feeling very agitated or restless                   | ◦ panic attacks                             |
| ◦ trouble sleeping (insomnia)                         | ◦ new or worse irritability                 |
| ◦ acting aggressive, being angry or violent           | ◦ acting on dangerous impulses              |
| ◦ an extreme increase in activity and talking (mania) | ◦ other unusual changes in behavior or mood |

**Do not take SPRAVATO<sup>®</sup> if you:**

- have blood vessel (aneurysmal vascular) disease (including in the brain, chest, abdominal aorta, arms and legs)
- have an abnormal connection between your veins and arteries (arteriovenous malformation)
- have a history of bleeding in the brain
- are allergic to esketamine, ketamine, or any of the other ingredients in SPRAVATO<sup>®</sup>.

If you are not sure if you have any of the above conditions, talk to your healthcare provider before taking SPRAVATO<sup>®</sup>.

**Before you take SPRAVATO<sup>®</sup>, tell your healthcare provider about all of your medical conditions, including if you:**

- have heart or brain problems, including:
  - high blood pressure (hypertension)
  - slow or fast heartbeats that cause shortness of breath, chest pain, lightheadedness, or fainting
  - history of heart attack
  - history of stroke
  - heart valve disease or heart failure
  - history of brain injury or any condition where there is increased pressure in the brain
- have liver problems
- have ever had a condition called “psychosis” (see, feel, or hear things that are not there, or believe in things that are not true).
- are pregnant or plan to become pregnant. SPRAVATO<sup>®</sup> may harm your unborn baby. You should not take SPRAVATO<sup>®</sup> if you are pregnant.

## IMPORTANT SAFETY INFORMATION (continued)

- Tell your healthcare provider right away if you become pregnant during treatment with SPRAVATO<sup>®</sup>.
- If you are able to become pregnant, talk to your healthcare provider about methods to prevent pregnancy during treatment with SPRAVATO<sup>®</sup>.
- There is a pregnancy registry for women who are exposed to SPRAVATO<sup>®</sup> during pregnancy. The purpose of the registry is to collect information about the health of women exposed to SPRAVATO<sup>®</sup> and their baby. If you become pregnant during treatment with SPRAVATO<sup>®</sup>, talk to your healthcare provider about registering with the National Pregnancy Registry for Antidepressants at 1-844-405-6185 or online at <https://womensmentalhealth.org/clinical-and-research-programs/pregnancyregistry/antidepressants/>.
- are breastfeeding or plan to breastfeed. SPRAVATO<sup>®</sup> passes into your breast milk. You should not breastfeed during treatment with SPRAVATO<sup>®</sup>.

**Tell your healthcare provider about all the medicines that you take**, including prescription and over-the-counter medicines, vitamins and herbal supplements. Taking SPRAVATO<sup>®</sup> with certain medicine may cause side effects.

**Especially tell your healthcare provider if you take** central nervous system (CNS) depressants, psychostimulants, or monoamine oxidase inhibitors (MAOIs) medicine. Keep a list of them to show to your healthcare provider and pharmacist when you get a new medicine.

### How will I take SPRAVATO<sup>®</sup>?

- You will take SPRAVATO<sup>®</sup> nasal spray yourself, under the supervision of a healthcare provider in a healthcare setting. Your healthcare provider will show you how to use the SPRAVATO<sup>®</sup> nasal spray device.
- Your healthcare provider will tell you how much SPRAVATO<sup>®</sup> you will take and when you will take it.
- Follow your SPRAVATO<sup>®</sup> treatment schedule exactly as your healthcare provider tells you to.
- During and after each use of the SPRAVATO<sup>®</sup> nasal spray device, you will be checked by a healthcare provider who will decide when you are ready to leave the healthcare setting.
- You will need to plan for a caregiver or family member to drive you home after taking SPRAVATO<sup>®</sup>.
- If you miss a SPRAVATO<sup>®</sup> treatment, your healthcare provider may change your dose and treatment schedule.
- Some people taking SPRAVATO<sup>®</sup> get nausea and vomiting. You should not eat for at least 2 hours before taking SPRAVATO<sup>®</sup> and not drink liquids at least 30 minutes before taking SPRAVATO<sup>®</sup>.
- If you take a nasal corticosteroid or nasal decongestant medicine take these medicines at least 1 hour before taking SPRAVATO<sup>®</sup>.

### What should I avoid while taking SPRAVATO<sup>®</sup>?

Do not drive, operate machinery, or do anything where you need to be completely alert after taking SPRAVATO<sup>®</sup>. Do not take part in these activities until the next day following a restful sleep. See **“What is the most important information I should know about SPRAVATO<sup>®</sup>?”**

### What are the possible side effects of SPRAVATO<sup>®</sup>?

#### SPRAVATO<sup>®</sup> may cause serious side effects including:

See **“What is the most important information I should know about SPRAVATO<sup>®</sup>?”**

**Increased blood pressure.** SPRAVATO<sup>®</sup> can cause a temporary increase in your blood pressure that may last for about 4 hours after taking a dose. Your healthcare provider will check your blood pressure before taking SPRAVATO<sup>®</sup> and for at least 2 hours after you take SPRAVATO<sup>®</sup>. Tell your healthcare provider right away if you get chest pain, shortness of breath, sudden severe headache, change in vision, or seizures after taking SPRAVATO<sup>®</sup>.

**Problems with thinking clearly.** Tell your healthcare provider if you have problems thinking or remembering.

**Bladder problems.** Tell your healthcare provider if you develop trouble urinating, such as a frequent or urgent need to urinate, pain when urinating, or urinating frequently at night.

#### The most common side effects of SPRAVATO<sup>®</sup> include:

- feeling disconnected from yourself, your thoughts, feelings and things around you
- dizziness
- nausea
- feeling sleepy
- spinning sensation
- decreased feeling of sensitivity (numbness)
- feeling anxious
- lack of energy
- increased blood pressure
- vomiting
- feeling drunk
- headache
- feeling very happy or excited

**If these common side effects occur, they usually happen right after taking SPRAVATO<sup>®</sup> and go away the same day.**

These are not all the possible side effects of SPRAVATO<sup>®</sup>.

Call your doctor for medical advice about side effects. You may report side effects to Johnson & Johnson at 1-800-526-7736, or to the FDA at 1-800-FDA-1088.

**Please see Important Safety Information throughout, full [Prescribing Information](#), including **Boxed WARNINGS**, and [Medication Guide](#) for SPRAVATO<sup>®</sup> and discuss any questions you may have with your healthcare provider.**

cp-170363v4

**Reference: 1.** Data on File. SD-217397. Janssen Pharmaceuticals, Inc.